

Permanent Rabi oscillations in coupled exciton-photon systems with PT-symmetry

I.Y. Chestnov,¹ S.S. Demirchyan,¹ S.M. Arakelian,¹ A.P. Alodjants,^{1,2} Y.G. Rubo,³ and A.V. Kavokin^{2,4}

¹*Department of Physics and Applied Mathematics,*

Vladimir State University named after A. G. and N. G. Stoletovs, 87 Gorkogo, Vladimir, 600000, Russia

²*Russian Quantum Center, 100 Novaya, Skolkovo, 143025, Moscow Region, Russia*

³*Instituto de Energías Renovables, Universidad Nacional Autónoma de México, Temizco, Morelos, 62580, Mexico*

⁴*Physics and Astronomy School, University of Southampton, Highfield, Southampton, SO171BJ, UK*

(Dated: March 26, 2015)

We propose a physical mechanism which enables permanent Rabi oscillations in driven-dissipative condensates of exciton-polaritons in semiconductor microcavities subjected to external magnetic fields. The method is based on incoherent excitonic reservoir engineering. We demonstrate that permanent non-decaying oscillations may appear due to the parity-time (PT) symmetry of the coupled exciton-photon system realised in a specific regime of pumping to the exciton state and depletion of the reservoir. For effective non-zero exciton-photon detuning, permanent Rabi oscillations occur with unequal amplitudes of exciton and photon components. Our predictions pave way to realisation of integrated circuits based on exciton-polariton condensates.

Introduction.—Nowadays, semiconductor microcavities with embedded quantum wells serve as a solid-state laboratory for fundamental studies of dynamic, coherent, nonlinear and quantum effects in nonequilibrium ensembles of bosonic quasiparticles: exciton-polaritons [1, 2]. Various approaches to photonic information processing with the use of microcavity polaritons have been proposed, see, e.g., [3–5]. In particular, polariton circuits, or neurons, are proposed in [6] and then experimentally examined by Ballarini *et al.* in [7] at extremely low pump intensities. Although nonequilibrium exciton-polariton Bose-Einstein condensation has been already observed in many labs, the requirement to use high Q-factor microcavities with high reflectivity Bragg mirrors limits wide application of such phenomena for practical purposes. Even in highest quality microcavities used in current experiments the leakage of photons leads to dissipation effects occurring on the scale of tens of picoseconds. In particular, this leads to a rapid attenuation of Rabi oscillations occurring between exciton and photon components of a polariton state, which represent a particular interest for quantum information applications. Recently, some of us have proposed [5] a way to improve the coherence time of polariton Rabi oscillations by stimulated pumping of a Rabi-oscillator from a permanent thermal reservoir of polaritons. The increase of the coherence time of polariton Rabi-oscillations in the presence of *cw* pumping has been experimentally observed [8], and manifestations of this effect in spatial dynamics of exciton-polaritons have been revealed experimentally [9] and analyzed theoretically [10]. Recently, Voronova *et al* [11] has studied theoretically the non-linear regime of polariton Rabi-oscillations and their interplay with exciton-photon Josephson oscillations.

In the Letter, we demonstrate theoretically realisation of permanent Rabi oscillations between excitonic and photonic components of a driven-dissipative spinor exciton-polariton condensate in semiconductor microcav-

ities at specific pumping conditions. We study the regime where the microcavity system obeys parity-time (PT) symmetry conditions. Originally, the PT-symmetry approach has been proposed by Bender *et al* [12, 13] to demonstrate that non-Hermitian Hamiltonians can possess entirely real energy eigenvalue spectra. In the past decade, PT-symmetry features have been demonstrated for various systems in photonics [14–19], condensed matter physics [20] and metamaterials [21]. Moreover, PT-symmetry approach has been extended to electronic circuits [22] which makes it highly relevant to polaritonics. Physically, PT-symmetry requirements can be easily understood by using the system of two linearly coupled oscillators (dimers, or waveguides in photonics [17, 18]). The system obeys PT-symmetry criteria if the rate of dissipation for one of the oscillators is exactly equal to the rate of gain in the other oscillator. This condition can be achieved in a system of two coupled polariton condensates with distributed dissipation rates [23], that results in the conservative Hamiltonian dynamics. It is important to note that even if the exact compensation between gain and losses in a two-mode system is not achieved, the concept of quasi-PT-symmetry can be introduced [24]. Here we demonstrate that the regime of permanent Rabi oscillations can be realized for the exciton-photon system in a microcavity, and the condition of PT-symmetry may be achieved if the gain in the excitonic component is compensated by losses from the photonic component.

Basic equations.—We consider a coupled exciton-photon system in the presence of the external magnetic field that produces the Zeeman splitting $\hbar\Delta_Z$ of exciton levels, and an incoherent excitonic reservoir, pumped by *cw*-field *P*. This non-resonant pumping can be realized both optically or by electronic current injection [25]. We assume that the wave function of driven-dissipative exciton-polariton condensate consists of photonic and excitonic components, $\phi_{\pm}$ and $\chi_{\pm}$. We will neglect the possible spatial degrees of freedom in what follows, assuming

the condensate to be at zero momentum state. In this case, both components of the condensate and the reservoir can be described by Boltzmann kinetic equations:

$$\dot{\chi}_{\pm} = \frac{1}{2} (p_X [N_{\pm}] - \gamma_X) \chi_{\pm} + i\delta_{\pm} \chi_{\pm} - i\Omega \phi_{\pm}, \quad (1a)$$

$$\dot{\phi}_{\pm} = -\frac{1}{2} \gamma_P \phi_{\pm} - i\Omega \chi_{\pm}, \quad (1b)$$

$$\dot{N}_{\pm} = P - \gamma_R N_{\pm} - p_X [N_{\pm}] |\chi_{\pm}|^2, \quad (1c)$$

where dots denote time derivatives and 2Ω is a polariton Rabi splitting frequency. The subscript “+” (“−”) corresponds to the spin projection parallel (antiparallel) to the vector of magnetic field. In Eqs. (1) $\delta_{\pm} = \Delta \pm \Delta_Z$ are effective photon-exciton detunings determined by the detuning $\Delta = \omega_P - \omega_X$ of the cavity mode and exciton frequencies in the absence of magnetic field and by the Zeeman splitting Δ_Z ; γ_X , γ_P and γ_R are the exciton, cavity and reservoir damping rates, respectively. In Eqs. (1) the term containing $p_X [N_{\pm}]$ describes the pumping of the exciton state by stimulated scattering from the reservoir. We shall consider two possible scattering mechanisms described by pumping terms p_1 and p_2 :

$$p_X [N_{\pm}] \equiv p_1 = R_1 N_{\pm}, \quad (2a)$$

$$p_X [N_{\pm}] \equiv p_2 = R_2 N_{\pm}^2 |\chi_{\pm}|^2. \quad (2b)$$

The Eq. (2a) implies the acoustic phonon assisted pumping with the rate R_1 , see [26], while Eq. (2b) describes the exciton-exciton scattering where excitons possessing momenta $-\mathbf{k}$ and $\mathbf{k}$ scatter into the condensate state with the momentum $\mathbf{k} = 0$. Note that in both cases, the scattering feeds both upper and lower exciton-polariton branches. Our model accounts for the decay of both exciton and photon components of the condensate but neglects incoherent processes that lead to relaxation between upper and lower polariton branches.

To start with, let us neglect scattering of excitons with opposite spins so that Eqs. (1) can be solved for each of spin components separately. We first examine the subsystem with “+” spin component omitting subscript for simplicity.

The dynamics of the exciton-polariton system demonstrates a threshold behavior at the specific value P^{th} of the cw-pump which is dependent on the particular mechanism of the exciton pumping $p_X [N]$ [26]. At $P < P^{th}$ the exciton pumping rate $p_X [N]$ is comparable to or much smaller than dumping rates γ_X and γ_P which, in turn, are much smaller than the Rabi splitting Ω (see, e.g., [27]). In this case one can assume the reservoir population N and the pumping rate $p_X [N]$ to be time independent on a time scale of the Rabi oscillation period $\sim \Omega^{-1}$. In particular, for the pumping term given by Eq. (2a), eliminating Eq. (1c) we find eigenfrequencies of

the exciton-photon system

$$\omega_{1,2} = \frac{i}{4} (\tilde{\gamma} - p_X) + \frac{\delta}{2} \pm \frac{1}{2} \sqrt{4\Omega^2 + \left(\delta - \frac{i}{2} (\gamma_P - \gamma_X + p_X) \right)^2}, \quad (3)$$

characterizing steady-state solutions $\chi = \chi_0 e^{i\omega t}$ and $\phi = \phi_0 e^{i\omega t}$. Physically, Eqs. (3) determine frequencies of upper (ω_1) and lower (ω_2) polariton branches, cf. [27]. In Eq. (3) we denoted $\tilde{\gamma} = \gamma_P + \gamma_X$. In this case, from Eq. (1c) one obtains $P_1^{th} = \tilde{\gamma} \gamma_R / R_1$.

The PT-symmetry of the system (1) appears if the hermitian part of Hamiltonian is symmetric and the anti-hermitian part is anti-symmetric with respect to permutation of excitonic and photonic components. The first condition leads to

$$\delta = 0, \quad (4a)$$

and the second condition requires that the losses from the photonic component should be fully compensated by the gain of the exciton one,

$$p_X - \gamma_X = \gamma_P. \quad (4b)$$

Clearly, if these conditions are satisfied the eigenfrequencies $\omega_{1,2}$ (3) become fully real. We note that in this case the amplitudes of oscillations of exciton and photon populations become equal $|\chi|^2 = |\phi|^2 = |A|^2$. Conditions (4a,b) can be experimentally realized by tuning of the pumping of the excitonic component with variation of the cw pumping strength.

Figure 1 reveals the decay dynamics of a coupled matter-light system determined by Eqs. (1) for different pumping mechanisms. The chosen parameters are relevant to experimentally accessible GaAs semiconductor microstructures, cf. [1]. In the Letter for simplicity we represented excitonic $|\chi|^2$ and photonic $|\phi|^2$ populations in the units of initial condensate density. As it is seen from Fig. 1 and follows from Eq. (3), the characteristic time $\tau_R \propto 2/(\tilde{\gamma} - p_X)$ of Rabi oscillations decay can be increased by manipulating the pump field parameter p_X .

Condition (4b) is a criterion for realisation of permanent Rabi oscillations ($\tau_R \rightarrow \infty$). It is easy to show that such a regime cannot be achieved below threshold, at $P < P^{th}$, see Fig. 1. Actually, since pumping rate $p_X \equiv p_X [N(t)]$ depends on the number of particles $N(t)$ in the reservoir, it varies as a function of time, and the condition (4b) just cannot be preserved in the whole time interval. This indicates that the features of the combined exciton-photon system are strongly affected by the reservoir dynamics, as the inset in Fig. 1 shows.

In the case of exciton pumping predominantly governed by the exciton-exciton scattering process Eq. (2b), the dynamics of the system is more complicated. The




FIG. 1. (Color online) Time dependence of the exciton population $|\chi|^2$ for $\delta = 0$ and for exciton state pumping p_1 (magenta curve) and p_2 (yellow curve) for cw-pump P below threshold P^{th} . Dash-dotted black curve indicates decay dynamics in the absence of reservoir. On the inset the relevant population of the reservoir N for exciton state pumping p_1 (magenta dashed) and p_2 (orange solid) is shown. Horizontal lines correspond to the stationary reservoir population, i.e. $N = P/\gamma_R$. Parameters are: $\gamma_X = 0.01 \text{ ps}^{-1}$, $\gamma_P = 0.1 \text{ ps}^{-1}$, $\hbar\Omega = 2.5 \text{ meV}$, $\gamma_R = 0.003 \text{ ps}^{-1}$, $\hbar R_1 = 0.01 \mu\text{m}^2\text{ps}^{-1}$ and $\hbar R_2 = 0.001 \mu\text{m}^2\text{ps}^{-1}$. Pumping rates are $P_1 = 0.02 \mu\text{m}^{-2}\text{ps}^{-1}$ and $P_2 = 0.035 \mu\text{m}^{-2}\text{ps}^{-1}$ – for magenta and yellow curves, respectively. Initial conditions are: $\chi(0) = 0$, $\phi(0) = 1$, $N(0) = P/\gamma_R$.

threshold value P_2^{th} can not be derived directly from Eq. (3) because of the dependence of p_2 term on the population $|\chi|^2$ of the excitonic component. Figure 1 (yellow curve) demonstrates the behaviour of exciton and photon components below threshold $P < P_2^{th}$, that was determined by numerical simulations. Although both components of the condensate are initially amplified, then the population of $|\chi|^2$ (and hence the pumping term $p_2 = R_2 N^2 |\chi|^2$) attenuate very rapidly.

Permanent oscillations.—Now we analyze permanent Rabi oscillations regime, that occurs above threshold $P > P^{th}$ and can be preserved as long as the PT-symmetry condition is fulfilled. We represent the solution of Eqs.(1a,b) in the form

$$\chi = \chi_1 e^{i\omega_1 t} + \chi_2 e^{i\omega_2 t}, \quad \phi = \phi_1 e^{i\omega_1 t} + \phi_2 e^{i\omega_2 t}, \quad (5a,b)$$

where $\chi_{1,2}$ and $\phi_{1,2}$ are constant amplitudes. In the permanent oscillations regime, the reservoir population oscillates with a small amplitude around some average value $\bar{N} = \langle N \rangle_t$. Further, we assume excitonic reservoir to be in the stationary state, $\dot{N} = 0$.

Let us first consider the phonon assisted pumping of the exciton component of the condensate. Substituting Eqs. (2a) and (5) into Eqs. (1a,b) and separating real and complex parts we obtain

$$R_1 \bar{N} = \gamma_X + \frac{\Omega^2 \gamma_P}{\gamma_P^2/4 + \omega_{1,2}^2}, \quad (6a)$$

$$\omega_{1,2}^3 - \delta \omega_{1,2}^2 - \omega_{1,2}(\Omega^2 - \gamma_P^2/4) - \delta \gamma_P^2/4 = 0. \quad (6b)$$

Clearly, only real roots of Eq. (6b) should be retained for the frequencies $\omega_{1,2}$.

Let us assume that the first requirement (4a) for the PT-symmetry is fulfilled, i.e., we set $\delta = 0$. In this case Eqs. (6) admit equal amplitudes $|A| \equiv |\chi_1| = |\chi_2| = |\phi_1| = |\phi_2|$ and characteristic polariton frequencies

$$\omega_{1,2} = \pm \omega \equiv \pm \sqrt{\Omega^2 - \gamma_P^2/4}. \quad (7)$$

Equation (7) is familiar in PT-symmetry theory for coupled oscillators, [15–17]. The value of $\Omega_c = \gamma_P/2$ can be associated with the PT-symmetry breaking threshold. In the case of a driven-dissipative exciton-polariton system, we operate significantly above the PT-threshold point, assuming that $\Omega \gg \Omega_c$, due to the strong exciton-photon coupling condition. Solution of Eq.(6a) with Eq.(7) leads to the condition

$$R_1 \bar{N} = \tilde{\gamma}, \quad (8)$$

that simply implies the balance between pump and loss rates in the system and follows directly from the PT-symmetry criterion (4b).

Figure 2 demonstrates the permanent oscillations regime for various pumping mechanisms of the excitonic component of the condensate. Permanent oscillations, shown in the left panel of the inset in Fig. 2, are established after several hundreds of picoseconds for experimentally accessible parameters. In particular, exciton and photon populations oscillate out of phase with equal constant amplitudes according to: $|\chi|^2 = 4|A|^2 \cos^2(\omega t)$ and $|\phi|^2 = 4|A|^2 \sin^2(\omega t)$, where $|A|^2 = (P - P_1^{th})/2\tilde{\gamma}$ can be determined from Eqs. (1c) and (8). Notably, the establishment of permanent Rabi oscillations regime is accompanied by the quick reservoir depletion (see the right panel on the inset in Fig. 2) resulting in the sharp peak of photon/exciton populations for the first few picoseconds – see Fig. 2. Undoubtedly, since the conditions (4) are satisfied now, this regime is also capable of having dynamical PT-symmetry properties occurring at the same time scale.

For $\delta \neq 0$ Eq. (6b) possesses three real roots, in general. Since the term $\delta \gamma_P^2/4$ might be small enough for the moderate values of the detuning δ we can omit this term and obtain (cf. (7))

$$\omega_{1,2} = \frac{1}{2} \left(\delta \pm \sqrt{\delta^2 + 4\Omega^2 - \gamma_P^2} \right). \quad (9)$$

The value of $\omega_R = |\omega_1 - \omega_2|$ represents the frequency of Rabi oscillations. However, since $|\omega_1| \neq |\omega_2|$, Eq. (6a) cannot be satisfied for both $\omega_{1,2}$ simultaneously, cf. (7). Hence, we can conclude that permanent oscillations cannot be supported in the case of $\delta \neq 0$, which also correlates with the first PT-symmetry criterion (4a).

Now let us examine the permanent oscillations regime for the exciton pumping by exciton-exciton scattering.




FIG. 2. (Color online) The same as in Fig. 1 but for cw-pump $P = 0.2 \mu\text{m}^{-2}\text{ps}^{-1}$, which is above threshold P^{th} . The relevant behavior of reservoir is shown on the right panel of the inset. Permanent Rabi oscillations for excitonic and photonic components, plotted on a zoomed time scale, are shown on the left panel of the inset.

In this case, using Eqs. (5) and omitting higher-order harmonics we obtain

$$R_2 \bar{N}^2 (|\chi_{1,2}|^2 + 2|\chi_{2,1}|^2) = \gamma_X + \frac{\Omega^2 \gamma_P}{\gamma_P^2/4 + \omega_{1,2}^2}. \quad (10)$$

Notably Eq. (6b) is valid in this case as well. Physically, it means that permanent Rabi oscillations frequency is independent on the physical mechanism of exciton pumping. The population $\bar{N}$ of the excitonic reservoir can be found from the equation

$$P - \gamma_R \bar{N} - R_2 \bar{N}^2 (|\chi_1|^4 + |\chi_2|^4 + 4|\chi_1|^2 |\chi_2|^2) = 0. \quad (11)$$

Again, in the limit of $\delta = 0$ we obtain a simple criterion $3R_2 \bar{N}^2 |A|^2 = \tilde{\gamma}$ of realization of permanent Rabi oscillations,

Figure 2 (yellow curve) demonstrates the population of excitonic $|\chi|^2$ component in this case. The frequency ω_R of oscillations is still determined by Eq. (7).

Now we consider permanent Rabi-oscillations in the case of non-zero detuning, $\delta \neq 0$, and exciton pumping p_2 . Equations (10) admit solutions simultaneously for both ω_1 and ω_2 in the case of unequal amplitudes, $\chi_1 \neq \chi_2$. These solutions exist only for the absolute value of detuning δ being smaller than some critical value δ_c . In fact, if $|\delta| > |\delta_c|$ population of an upper or a lower polariton branch (depending on the sign of δ) at the steady state becomes zero, that indicates the collapse of Rabi oscillations. It means that the initial system (1) has a steady state solution only; exciton $|\chi|^2$ and photon $|\phi|^2$ populations tend to some constant (non-zero) values determined by this solution. The value of δ_c can be obtained using Eqs. (9) and (10) and it approximately reads

$$\delta_c \simeq \pm \left(\frac{\Omega^2}{4\gamma_P} \left[\gamma_X - 8\gamma_P + 3\sqrt{\gamma_X^2 + 8\gamma_P^2} \right] \right)^{1/2}. \quad (12)$$




FIG. 3. Time evolution of Stokes parameters for $\Delta_Z = 0.2\Omega$, $\Delta = 0.15\Omega$. S_x (red) and S_y (blue) components – (a); S_z (red) and S_0 (blue) component – (b); (c) and (d) show the evolution of polarization on (S_x, S_y) -plane (c) and on Stokes sphere (d) on the time scale of $T_B = 2\pi/\omega_B$ characterizing permanent Rabi oscillations beating period. The parameters are: $p_X[N] = p_2$, $P = 0.2 \mu\text{m}^{-2}\text{ps}^{-1}$; other values are the same as in Fig. 1.

It is important that δ_c is determined by the parameters of a Rabi-oscillator as a whole, i.e., by the Rabi splitting frequency and damping rates γ_X , γ_P , and it does not depend on the parameters of the reservoir, including the external pump P . For GaAs-based semiconductor microcavities we obtain $\hbar\delta_c \simeq 0.38\hbar\Omega = 0.96 \text{ meV}$. Thus, permanent oscillations may be established in the vicinity of exciton-photon resonance for experimentally accessible parameters.

Polarization properties.—Here we consider a complete set of Eqs. (1), that characterizes spin dependent exciton-polariton properties. Polarization properties of the excitonic system are conveniently described by the Stokes vector $\mathbf{S}$ with components $S_{x,y,z}$ defined by $\mathbf{S} = \frac{1}{2}(\Psi^\dagger \cdot \boldsymbol{\sigma} \cdot \Psi)$, where $\Psi = (\chi_+, \chi_-)^T$ and $\sigma_{x,y,z}$ are the Pauli matrices. Since $\delta_+ \neq \delta_-$ the behavior of polarization of exciton component is completely determined by detuning Δ for a given non-zero Zeeman shift $\Delta_Z \neq 0$.

The general case corresponding to both Δ and Δ_Z are nonzero is illustrated by Fig. 3. Using anzats (5) and definition for Stokes vector $\mathbf{S}$ it is easy to show that components S_x and S_y oscillate with the frequency Δ_Z (frequent beats in Figs. 3a and 4a). Besides there are fast oscillations with the frequency $(\omega_{R+} + \omega_{R-})/2$ close to the frequency of Rabi-oscillations. If the conditions of establishing of permanent oscillations are fulfilled for both spin components simultaneously ($|\delta_\pm| < \delta_c$), the frequencies of Rabi oscillations ω_{R+} and ω_{R-} are dif-




FIG. 4. Time evolution of Stokes parameters $S_{x,y,z}$ for $\Delta_Z = \Delta = 0.2\Omega$. The parameters are: $p_X[N] = p_1$, $P = 0.2 \mu\text{m}^{-2}\text{ps}^{-1}$; other values are the same as in Fig. 1

ferent, because of the relation $|\delta_+| \neq |\delta_-|$. In this case, the amplitudes of Stokes vector oscillations undergo additional long-period beats with the period $T_B = 2\pi/\omega_B = 2\pi/(\omega_{R+} - \omega_{R-}) \approx \frac{2\pi\Omega}{\Delta\Delta_Z}$, see Fig. 3(b). These beats are sustained in time.

In Fig. 3(d) we represented the evolution of the Stokes vector $\mathbf{S}$ normalised to the total number of excitons $S_0 = \frac{1}{2}(|\chi_+|^2 + |\chi_-|^2)$ on the Poincare sphere with a unit radius. Note that S_0 , describing the excitonic subsystem, oscillates in time as shown in Fig. 3(b).

Let us consider the limit where permanent oscillations disappear for one of the spin components, i.e. for $|\delta_+| < |\delta_c|$ and $|\delta_-| > |\delta_c|$, or vice versa, that can be achieved by the appropriate choice of Δ_Z and Δ . In fact, in this case the beats in Stokes parameters with the period T_B are suppressed in time and the amplitude of oscillations tends to the constant value. Figure 4 demonstrates these features for Stokes parameters in the specific case of $\Delta = \Delta_Z$. In this limit, the detunings $\delta_- = 0$ and $\delta_+ = 2\Delta_Z$. Physically, it means that permanent oscillations can be established for an exciton pumping mechanism described by (2a), but only for “-” spin component while population of another component will approach a constant value. At long time intervals only fast oscillations survive.

In conclusion, we have described the regime of permanent Rabi oscillations in a driven-dissipative exciton-photon system feeded from the incoherent exciton reservoir. Two types of reservoir-Rabi-oscillator coupling mechanisms have been examined. We have shown that the permanent Rabi oscillation regime is established above the threshold pumping where the dynamical PT-symmetry of the coupled exciton-photon system is realised. Notably, this regime occurs in a non-resonant case only if the exciton component of the condensate is pumped by exciton-exciton scattering predominantly. We studied excitonic polarization properties in the presence of external magnetic fields. The conditions for permanent Rabi oscillations for polarized excitonic and/or photonic system are established and discussed. Realization of permanent Rabi oscillations in this case play a crucial role for creation and manipulating of long-lived spin polarisation in the exciton-photon system which pave the way to the design of new optical memory devices.

Acknowledgments.—A.P.A. acknowledges useful discussions with Yuri Kivshar. This work was supported by RFBR Grants No. 14-02-31443, No. 14-02-92604, No. 14-02-97503, No. 15-52-52001, No. 15-59-30406, No. 14-32-50438 and EU project PIRSESGA-2013-612600 LIMACONA. The financial support from the Russian Ministry of Education and Science (Contract No.11.G34.31.0067 with SPbSU and leading scientist A. V. K.) is acknowledged. A.P.A. acknowledges support from “Dynasty” Foundation and Y.G.R. acknowledges support from CONACYT under the grant No. 251808.

-
- [1] H. Deng, H. Haug, Y. Yamamoto, *Rev. Mod. Phys.* **82**, 1489 (2010).
 - [2] *Phys. Status Solidi B* **242**(11) (2005), special issue of Physics of Semiconductor Microcavities, edited by B. Deveaud.
 - [3] C. Ciuti, *Phys. Rev. B* **69**, 245304 (2004); M. Romanelli *et al.*, *Phys. Rev. B* **82**, 155313 (2010).
 - [4] D. Pagel, H. Fehske, J. Sperling, and W. Vogel, *Phys. Rev. A* **86**, 052313 (2012); S. Portolan *et al.*, *New J. Phys.* **16**, 063030 (2014).
 - [5] S.S. Demirchyan *et al.*, *Phys. Rev. Lett.* **112**, 196403 (2014).
 - [6] C. Leyder *et al.*, *Phys. Rev. Lett.* **99**, 196402 (2007); T.C.H. Liew, A.V. Kavokin, and I.A. Shelykh, *ibid.* **101**, 016402 (2008).
 - [7] D. Ballarini *et al.*, *Nat. Commun.* **4**, 1778 (2013).
 - [8] M. De Giorgi *et al.*, *Phys. Rev. Lett.* **112**, 113602 (2014).
 - [9] L. Dominici *et al.*, *Phys. Rev. Lett.* **113**, 226401 (2014).
 - [10] T.C.H. Liew, Y.G. Rubo, and A.V. Kavokin, *Phys. Rev. B* **90**, 245309 (2014).
 - [11] N.S. Voronova, A.A. Elistratov and Yu.E. Lozovik, *arXiv:1503.04231* (2015).
 - [12] C. M. Bender and S. Boettcher, *Phys. Rev. Lett.* **80**, 5243 (1998).
 - [13] C. M. Bender, S. Boettcher, and P. N. Meisinger, *J. Math. Phys. (N.Y.)* **40**, 2201 (1999).
 - [14] C.E. Rüter *et al.*, *Nat. Phys.* **6**, 192 (2010).
 - [15] L. Feng, M. Ayache, J. Huang, *et al.*, *Science* **729**, 333 (2011).
 - [16] A. Szameit, M. C. Rechtsman, O. Bahat-Treidel, and M. Segev, *Phys. Rev. A* **84**, 021806(R) (2011).
 - [17] H. Ramezani and T. Kottos, R. El-Ganainy and D. N. Christodoulides, *Phys. Rev. A* **82**, 043803 (2010).
 - [18] A.A. Sukhorukov, Z. Xu, and Y.S. Kivshar, *Phys. Rev. A* **82**, 043818 (2010).
 - [19] I.V. Barashenkov, S.V. Suchkov, A.A. Sukhorukov, S.V. Dmitriev, and Y.S. Kivshar, *Phys. Rev. A* **86**, 053809 (2012).
 - [20] E.-M. Graefe, *J. of Phys. A, Math. And Theor.* **45**, 444015 (2012); C. Hang, G. Huang, and V.V. Konotop, *Phys. Rev. Letts.* **110**, 083604 (2013).
 - [21] G. Castaldi *et al.*, *Phys. Rev. Lett.* **110**, 173901 (2013).
 - [22] J. Schindler *et al.*, *Math. and Theor.* **45**, 444029 (2012).
 - [23] I.L. Aleiner, B.L. Altshuler, and Y.G. Rubo, *Phys. Rev. B* **85**, 121301(R) (2012).
 - [24] M. Ornigotti and A. Szameit, *J. Opt.* **16**, 065501 (2014).
 - [25] C. Schneider, A. Rahimi-Iman, N.Y. Kim, *et al.*, *Nature*

- 497**, 348 (2013); P. Bhattacharya, Bo Xiao, A. Das, S. Bhowmick, and J. Heo, Phys. Rev. Lett. **110**, 206403 (2013).
- [26] M. Wouters, I. Carusotto, Phys. Rev. Lett. **99**, 140402 (2007).
- [27] V. Savona, L.C. Andreani, P. Schwendimann, and A. Quattropani, Solid State Comms **93**, 733 (1995).